

A Cosmographical Approach to the Redshift Drift in $f(R)$ Theories

Florencia Anabella Teppa Pannia*

*Facultad de Ciencias Astronómicas y Geofísicas, Universidad Nacional de La Plata,
Paseo del Bosque s/n, B1900FWA, Argentina.*

Santiago Esteban Perez Bergliaffa†

*Departamento de Física Teórica, Instituto de Física,
Universidade do Estado de Rio de Janeiro, CEP 20550-013, Rio de Janeiro, Brazil.*

By means of a cosmographical approach to the redshift drift, together with a series development of the Hubble parameter for an arbitrary $f(R)$ theory, we obtain a restriction on f and some of its derivatives evaluated at the present epoch. It is show that, when applied to specific forms of $f(R)$, this restriction (which depends on the values of the kinematical parameters today) leads to bounds on the parameters of the theory that are more stringent than those previously obtained.

PACS numbers: 04.50.Kd, 98.80.Es

Keywords: cosmography; modified theories of gravity

I. INTRODUCTION

The interpretation of several sets of data (such as those obtained from type Ia supernovae, large scale structure, baryon acoustic oscillations, and the cosmic microwave background) in the framework of the Standard Cosmological Model (SCM) (based on General Relativity (GR) and the Cosmological Principle (CP)) indicates that the universe is currently undergoing a phase of accelerated expansion. The most commonly accepted candidates to source such an expansion are the cosmological constant and some unknown type of matter dubbed “dark energy” (for a complete list see [1]). While the energy density associated with the cosmological constant that is inferred from astronomical observations is approximately 120 orders of magnitude lower than the value predicted by field theory (see for instance [2]), the scalar field used to model dark energy has features that are alien to those displayed by the scalar fields of particle physics [3]. An alternative to describe the accelerated expansion is to assume that a theory different from GR governs the dynamics of the gravitational field for very low values of the curvature. The simplest generalization of the Einstein-Hilbert Lagrangian is given by

$$S = \int \sqrt{-g} f(R) d^4x,$$

where the dependence of the function f on the scalar curvature R is to be determined by several criteria (such as matter stability [4], absence of ghost modes in the cosmological perturbations [5], and correct succession of cosmological eras [6]), and by the comparison of the predictions of $f(R)$ theories with observation¹ (for instance, the stability of cosmological perturbations [9]). Several forms for $f(R)$ have been constructed in order to successfully satisfy these constraints (for instance those given in [10, 11]).

Currently a lot of effort is devoted to the study of quantities and effects that have the potential of discriminating between different $f(R)$ models, and between these models and the SCM. Among them, we can mention the growth rate of matter density perturbations (see for instance [12]), the enhanced brightness of dwarf galaxies [13], the modifications of the 21cm power spectrum at reionisation [14], the specific angular momentum of galactic halos [15], and the number

*Electronic address: fteppa@fcaglp.unlp.edu.ar

†Electronic address: sepbergliaffa@gmail.com

¹ For reviews about different aspects of $f(R)$ theories, see [2, 3, 7, 8].

counts of peaks in weak lensing maps [16]. We shall show that another observable that can be of use along these lines is the redshift drift ($\Delta z/\Delta t$), that is, the time variation of the cosmological redshift caused by the expansion of the Universe. The redshift drift was first considered by Sandage [17], and the effects of a nonzero cosmological constant on it were presented in [18]. In 1998, Loeb [19] discussed the possibility of actually measuring the redshift drift of cosmological sources using spectroscopic techniques available at that time. Since then, the potential of planned extremely large observatories, such as the European Extremely Large Telescope (E-ELT), the Thirty Meter Telescope and the Giant Magellan Telescope, and the development of ultra-stable spectrographs with high-resolution such as the COsmic Dynamics and EXo-earth experiment (CODEX) proposed for the E-ELT, together with simulations, showed that the measurement of the redshift drift is feasible in the near future (for a review, see [20]), leading to what has been dubbed “real-time cosmology” in [21].

When compared to other cosmological observables, the redshift drift has the advantage that it directly tests the dependence of the Hubble parameter with the redshift, hence probing the dynamics of the scale factor. Another feature of this observable is that it does not depend on details of the source (such as the absolute luminosity), or on the definition of a standard ruler. As demonstrated in [22], the measurement of the redshift drift would allow the test of the Copernican Principle, thus checking for any degree of radial inhomogeneity. This issue was further discussed in [23], where it was shown that the redshift drift is positive for sources with $z < 2$ in the Λ CDM, while in Lemâitre-Tolman-Bondi models (see for instance [24]) is negative for sources observed from the symmetry center [25]. The redshift drift can also be used to constrain phenomenological parametrizations of dynamical dark energy models (see [26–28]). In this work, we shall extend the applications of the redshift drift by showing that it can be employed to set limits on the parameters of an arbitrary $f(R)$ theory. This goal will be achieved by combining a cosmographical approach to the redshift drift with the corresponding dynamics.

Cosmography is a mathematical framework for the description of the universe, based entirely on the CP, and on those parts of GR that follow directly from the Principle of Equivalence [29]. It is inherently kinematic, in the sense that it is independent of the dynamics obeyed by the scale factor $a(t)$. The cosmographical approach has already led to interesting results in the framework of $f(R)$ theories [30–33]. It will be used in Sect. II to obtain a series expansion of the redshift drift in terms of the cosmological redshift z and the derivatives of the scale factor. Since the redshift drift depends on the explicit form of the Hubble parameter $H(z)$, we shall show in Sect. III that its series expansion can actually be written in terms of $f(R)$ and its derivatives, using the Equations of Motion (EoM) for the $f(R)$ theory in the metric version. By comparing the two series, we show that there exists relations between $f(R)$, its derivatives, and the kinematical parameters, which impose constraints on the parameters of a given $f(R)$. By using one of these relations in two examples: that proposed by Hu and Sawicki [11] and the exponential gravity theory introduced by Linder [34]. In both cases, we find new limits for the parameter space of each theory and compare our results with previous constraints. We close in Sect. IV with some remarks.

II. A COSMOGRAPHICAL APPROACH TO THE REDSHIFT DRIFT

In this section we shall sketch the calculation that leads to the expression for the series expansion of the redshift drift in terms of z in homogeneous and isotropic space-times. The redshift of a photon emitted by a source at $t = t_s$ that reaches the observer at $t = t_{obs}$ is given by

$$z(t_s) = \frac{a(t_{obs})}{a(t_s)} - 1. \quad (1)$$

The time variation of the redshift is obtained by comparing this expression with the one corresponding to a photon emitted at $t = t_s + \Delta t_s$, that is $\Delta z = z(t_s + \Delta t_s) - z(t_s)$. To first order in Δt_{obs} and Δt_s , it follows that [19]

$$\frac{\Delta z}{\Delta t_{obs}} = \left[\frac{\dot{a}(t_{obs}) - \dot{a}(t_s)}{a(t_s)} \right]. \quad (2)$$

Using the definition of z and $H(z)$ in this equation we get the expression of the redshift drift in terms of z , namely

$$\frac{\Delta z}{\Delta t_{obs}} = (1+z)H_{obs} - H(z). \quad (3)$$

In the case of the SCM, the redshift drift can be written as a function of the cosmological parameters using the corresponding expression for $H(z)$ as follows [19]:

$$(\Delta z/\Delta t) = H_0 \left[(1+z) - (\Omega_{m,0}(1+z)^3 + \Omega_{r,0}(1+z)^4 + \Omega_{\Lambda,0})^{1/2} \right], \quad (4)$$

where the subindex 0 means that the corresponding quantity is evaluated at $t = \text{today}$. As discussed in the Introduction, we would like to pursue here a mixed approach partially based in the concept of cosmography (see for instance [35]). Cosmography consists in constraining the history of the scale factor (and a posteriori, of quantities defined in terms of $a(t)$) from observational data, without using the dynamics obeyed by the gravitational field. The series development of the scale factor around t_0 is given by

$$a(t) = a_0 \left[1 + H_0(t-t_0) - \frac{1}{2}q_0 H_0^2(t-t_0)^2 + \frac{1}{3!}j_0 H_0^3(t-t_0)^3 + \frac{1}{4!}s_0 H_0^4(t-t_0)^4 + \mathcal{O}[(t-t_0)^5] \right], \quad (5)$$

where the so-called kinematical parameters are defined by

$$H(t) \equiv +\frac{1}{a(t)} \frac{da}{dt}; \quad q(t) \equiv -\frac{1}{a(t)} \frac{d^2 a}{dt^2} \left[\frac{1}{a(t)} \frac{da}{dt} \right]^{-2}; \quad j(t) \equiv +\frac{1}{a(t)} \frac{d^3 a}{dt^3} \left[\frac{1}{a(t)} \frac{da}{dt} \right]^{-3}; \quad s(t) \equiv +\frac{1}{a(t)} \frac{d^4 a}{dt^4} \left[\frac{1}{a(t)} \frac{da}{dt} \right]^{-4}.$$

In order to use Eq. (5) for the calculation of the redshift drift, we need to express t_s in terms of known quantities. This can be done through the physical distance travelled by a photon emitted at t_s and observed at t_0 , given by

$$D = c \int dt = c(t_0 - t_s). \quad (6)$$

This equation, together with (1) yields

$$z(D) = \frac{H_0 D}{c} + \frac{2+q_0}{2} \frac{H_0^2 D^2}{c^2} + \frac{6(1+q_0)+j_0}{6} \frac{H_0^3 D^3}{c^3} + \mathcal{O} \left(\left[\frac{H_0 D}{c} \right]^4 \right), \quad (7)$$

which can be inverted to [35]

$$D(z) = \frac{c z}{H_0} \left[1 - \left(1 + \frac{q_0}{2} \right) z + \left(1 + q_0 + \frac{q_0^2}{2} - \frac{j_0}{6} \right) z^2 - \left(1 + \frac{3}{2} q_0(1+q_0) + \frac{5}{8} q_0^3 - \frac{1}{2} j_0 - \frac{5}{12} q_0 j_0 - \frac{s_0}{24} \right) z^3 + \mathcal{O}(z^4) \right]. \quad (8)$$

Using Eqs. (6) and (8) we can write $a(t_s)$ as a power series in z , with coefficients that are functions of the kinematical parameters. After some algebra, it follows that

$$\frac{\Delta z}{\Delta t_0} = -H_0 q_0 z + \frac{1}{2} H_0 (q_0^2 - j_0) z^2 + \frac{1}{2} H_0 \left[\frac{1}{3} (s_0 + 4q_0 j_0) + j_0 - q_0^2 - q_0^3 \right] z^3 + \mathcal{O}(z^4). \quad (9)$$

This equation gives the expression of the redshift drift up to the third order in the redshift of the source, in terms of the value of the kinematical parameters at the present epoch. Note that the coefficient of z^n involves the $(n+1)$ -derivative of the scale factor.

In the next section we shall obtain an expression for the redshift drift using the dynamics of an arbitrary $f(R)$ theory together with the development in terms of z .

III. THE REDSHIFT DRIFT IN $f(R)$ THEORIES

Let us recall that the redshift drift can be expressed in terms of $H(z)$ as follows

$$\frac{\Delta z}{\Delta t_0} = (1+z)H_0 - H(z) \quad (10)$$

The dynamics of a given theory establishes the form of the function $H(z)$. In the case of the SCM, we obtain for the redshift drift the expression given in Eq. (4) using Friedmann equation. In the case of $f(R)$ theories, the EoM that follow from the variation of the action

$$S = \int d^4x \sqrt{-g} [f(R) + \mathcal{L}_{\text{matter}}]$$

w.r.t. the metric must be used instead. For the FLRW metric and considering a pressureless cosmological fluid, these equations are (see for instance, [36])

$$f'R - 2f - 3f'' \left(\ddot{R} + \frac{3\dot{a}\dot{R}}{a} \right) - 3f''' \dot{R}^2 = T, \quad (11)$$

$$f'R_{tt} + \frac{1}{2}f + 3f'' \frac{\dot{a}\dot{R}}{a} = T_{tt}, \quad (12)$$

where dot and prime denote, respectively, derivative w.r.t. t and R , $R_{tt} = 3\ddot{a}/a$, $R = 6(\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2})$, and T is the trace of the energy-momentum tensor. From these, an equation for H can be obtained, and it reads

$$H = -\frac{1}{6\dot{R}f''} (6H^2f' - 2\rho_m - f + Rf'). \quad (13)$$

Replacing Eq. (10), we obtain

$$\frac{\Delta z}{\Delta t_0}(z) = \frac{\dot{a}_0 - \dot{a}(z)}{a(z)} = (z+1)H_0 + \frac{1}{6} \frac{f(R) - Rf'(R) + 6H^2f'(R) - 2\rho_m}{\dot{R}(t)f''(R)}, \quad (14)$$

where R , H , and ρ_m are functions of z . Expanding this equation in powers of z and using the definitions of the kinematical parameters, the redshift drift up to first order in z for an arbitrary $f(R)$ is given by

$$\begin{aligned} \frac{\Delta z}{\Delta t_0}(z) = & \{ [j_0^2(1-q_0) + 4(1-j_0) + 2q_0j_0(1+q_0) + q_0^2(3-q_0)] 36H_0^4(f_0'')^2 \\ & + [(s_0 + q_0^2 + 6 + 8q_0)(f_0 - 2\rho_m) + (2j_0(1+q_0) + q_0(s_0 + q_0^2 + 6q_0) - 4) 6H_0^2f_0'] f_0'' \\ & + [6q_0H_0^2f_0' - 2\rho_m + f_0] [(q_0 - j_0)^2 + 4(1+q_0 - j_0)] 6H_0^2f_0''' \} \frac{z}{36(f_0'')^2(j_0 - q_0 - 2)^2H_0^3} + \mathcal{O}(z^2). \end{aligned} \quad (15)$$

where we have used the expressions $R = 6(\dot{H} + 2H^2)$, $\dot{H} = -H^2(1+q)$, and $\ddot{H} = H^3(j+3q+2)$. The second order term involves derivatives of the scale factor higher than the fourth.

Comparing expression (15) with the linear term of Eq. (9) we find a relation between $f(R)$, its derivatives and the kinematical parameters, all evaluated at $t = \text{today}$:

$$\begin{aligned} & [(s_0 + q_0^2 + 6 + 8q_0)(f_0 - 2\rho_m) + (-4 + 6q_0^2 + 2j_0 + q_0s_0 + 2j_0q_0 + q_0^3)6H_0^2f_0'] f_0'' + \\ & + [(q_0 - j_0)^2 + 4(1+q_0 - j_0)] [(f_0 - 2\rho_m + 6q_0H_0^2f_0')f_0''' + 6H_0^2(f_0'')^2] 6H_0^2 = 0. \end{aligned} \quad (16)$$

Since we are working at first order in z , this equation is a necessary condition for any $f(R)$ theory to describe the variation of the redshift drift with z . By equating higher orders of z from Eqs. (9) and (15) we would obtain more (actually, an infinite number of) necessary conditions on $f(R)$ and its derivatives. If the theory under discussion is to describe the redshift drift at all orders in z , all these conditions should be satisfied. Let us remark that the condition that would be obtained from the second order in z would not be restrictive, since it will depend on the value of the kinematical parameter ℓ_0 (proportional to the fifth derivative of the scale factor), for which there are no observational limits available.

We shall see next how Eq. (16) constrains the value of the parameters of a given $f(R)$, by applying it to two particular forms of $f(R)$.

A. Example 1: The Theory of Hu and Sawicki

Let us start with the theory introduced by Hu and Sawicki [11], which is given by

$$f(R) = R - m^2 \frac{c_1(R/m^2)^n}{c_2(R/m^2)^n + 1}, \quad (17)$$

where $n > 0$, c_1 and c_2 are dimensionless parameters and the mass scale is $m^2 \equiv \kappa^2 \bar{\rho}_0/3$, with $\bar{\rho}_0$ the average density today.

For values of the curvature high compared with m^2 , $f(R)$ may be expanded as

$$\lim_{m^2/R \rightarrow 0} f(R) \simeq R - \frac{c_1}{c_2} m^2 + \frac{c_1}{c_2^2} m^2 (m^2/R)^n \quad (18)$$

and, at finite c_1/c_2^2 , the theory can approximate the expansion history of the Λ CDM model [11]. In this regime, the parameters c_1 and c_2 must satisfy the relation

$$\frac{c_1}{c_2} \approx 6 \frac{\tilde{\Omega}_\Lambda}{\tilde{\Omega}_m}. \quad (19)$$

For the flat Λ CDM expansion history, Eq. (18) yields

$$R \simeq 3m^2 \left(a^{-3} + 4 \frac{\tilde{\Omega}_\Lambda}{\tilde{\Omega}_m} \right), \quad (20)$$

$$f' \simeq 1 - n \frac{c_1}{c_2^2} \left(\frac{m^2}{R} \right)^{n+1}, \quad (21)$$

and at the present epoch,

$$R_0 \simeq m^2 \left(\frac{12}{\tilde{\Omega}_{m,0}} - 9 \right), \quad (22)$$

$$f'_0 \simeq 1 - n \frac{c_1}{c_2^2} \left(\frac{12}{\tilde{\Omega}_{m,0}} - 9 \right)^{-n-1}. \quad (23)$$

Using Eq. (23), c_1/c_2^2 can be expressed in terms of n , f'_0 and $\tilde{\Omega}_{m,0}$. In addition, higher order derivatives of f can also be written in terms of the same quantities. Hence from now on we set $\tilde{\Omega}_{m,0} = 0.274 \pm 0.007$ [37] and leave f'_0 and n the only free parameters of the **theory**. With these considerations, Eq. (16) yields

$$f'_0(n) = \frac{1}{A} \left\{ [q_0(8 + 5q_0 - 6j_0 + (q_0 - j_0)^2) + (j_0 - 2)^2] 4n + [j_0(j_0 - 2q_0 - 4) - 14 - 2q_0(10 + q_0) - 3s_0] 6\tilde{\Omega}_{m,0} \right. \\ \left. + [q_0(8 + q_0) + 6 + s_0] 9\tilde{\Omega}_{m,0}^2 + 4 [2(q_0(10 - 2j_0 + 3q_0) + 6 + s_0) + q_0(q_0 - j_0)^2] \right\}, \quad (24)$$

with

$$A = [q_0(8 + 5q_0 - 6j_0 + (q_0 - j_0)^2) + (j_0 - 2)^2] 4n - [2q_0(13 + 2q_0) - j_0(j_0 - 2) + 18 + 3s_0] 6\tilde{\Omega}_{m,0} \\ + [q_0(8 + q_0) + 6 + s_0] 9\tilde{\Omega}_{m,0}^2 + 4 [2(10 + 5q_0^2 + 16q_0 - 2j_0(2q_0 + 1) + s_0) + q_0(q_0 - j_0)^2].$$

Expression (24) gives a relation between the parameters f'_0 and n in terms of $\tilde{\Omega}_{m,0}$ and the kinematical parameters H , q , j , and s , all evaluated today. We shall take the values $H_0 = 74.2 \pm 3.6$ km/s/Mpc [38], $q_0 = -0.669 \pm 0.052$, $j_0 = 0.284 \pm 0.151$ and $s_0 = -0.680 \pm 0.456$ [31]. In Figure 1 we plot the relation $f'_0(n)$ provided by our cosmographical approach to the redshift drift combined with the expansion of the expression of $H(z)$ for $f(R)$ theories. The curve tends asymptotically to $f'_0 = 1$, which corresponds to the GR limit. The plot also displays the limit obtained from solar system tests, given by $|f'_0 - 1| < 0.1$ [11]. We find that actually $1 - f'_0 < 0.1$ and, from this limit, values for n larger than 12.8 are favoured. This result improves the previous bound for the parameter n obtained from measurements of the linear growth of structure and SNIa luminosity [39].




FIG. 1: Constraints on the parameter space of the theory by Hu and Sawicki. The red line corresponds to the relation between f'_0 and n that follows from Eq. (16). The dashed lines represent the error propagation arising from the kinematical parameters and the density matter at the present epoch. The shady region indicates the values of f'_0 in agreement with solar system tests.

B. Example 2: Exponential Gravity

Next we shall analyze the restrictions that follow from Eq. (16) on the choice of $f(R)$ proposed by Linder [34]:

$$f(R) = R - cr(1 - \exp(-R/r)), \quad (25)$$

where c and r are two (positive) parameters of the model. This $f(R)$ was specifically designed to (i) avoid the inclusion of an implicit cosmological constant (since it vanishes in the low curvature limit), (ii) reduce to GR for high values of the curvature, (iii) incorporate a transition scale (given by r) to be fitted from observations (instead of set equal to R_0), and (iv) restore GR for locally high curvature systems such as the solar system or galaxies. As shown in [34], the product cr is given in terms of Ω_m by

$$cr = 6m^2(\Omega_m^{-1} - 1). \quad (26)$$

The use of Eq. (16) for the current choice of $f(R)$ yields a relation between the dimensionless parameters c and r/m^2 given by

$$c \left(\frac{r}{m^2} \right) = \frac{1}{B} \left\{ -6 \frac{r}{m^2} \Omega_m [(q_0(8 + q_0) + 6 + s_0) \Omega_m - (2(1 + j_0 q_0 + q_0 + j_0) + s_0 - q_0^2)] \right. \\ \left. + 36 [4(q_0 + 1 - j_0) + (q_0 - j_0)^2] (\Omega_m + 1) \right\} \exp \left(\frac{6(1 - q_0)}{\frac{r}{m^2} \Omega_m} \right), \quad (27)$$

with

$$B = \left\{ [q_0(8 + q_0) + 6 + s_0] \left[\exp \left(\frac{6(1 - q_0)}{\frac{r}{m^2} \Omega_m} \right) - 1 \right] \right\} \left(\frac{r}{m^2} \Omega_m \right)^2 \\ + 6 \left\{ q_0(q_0^2 + 7q_0 + 4 + s_0) + j_0(j_0 - 2) - [4(q_0 + 1 - j_0) + (q_0 - j_0)^2] \exp \left(\frac{6(1 - q_0)}{\frac{r}{m^2} \Omega_m} \right) \right\} \frac{r}{m^2} \Omega_m \\ - 36 [4(1 + 2q_0) + q_0^2(5 + q_0) + j_0^2(1 + q_0) - j_0(2q_0^2 + 6q_0 + 4)],$$

which is plotted along with Eq. (26) in Figure 2. The strips formed by both curves and the corresponding errors juxtapose for all values of $c \gtrsim 12$, which imply that $r/m^2 \lesssim 1.4$. Notice that the range of possible values for the parameter c that follows from our method considerably improves the previous bound ($c \geq 1.27$) obtained in [40].




FIG. 2: Constraints coming from our approach to the redshift drift (continuous line) and the relation (26) between the parameters of the theory (dashed line). In both cases, the thin lines represent the error propagation arising from the kinematical parameters and the matter density at the present epoch.

IV. DISCUSSION

We have developed a cosmographical approach to the redshift drift and compared it with a series expansion in terms of z of the Hubble parameter for an arbitrary $f(R)$ theory. As a result, a relation between $f(R)$, its derivatives and the value of the kinematical parameters, all evaluated at the present epoch, was obtained. This relation must be satisfied for a given $f(R)$ theory to describe the redshift drift at first order in z . The comparison of higher-order terms would yield relations between the quantities mentioned above, that involve kinematical parameters of order higher than the fourth (which have not been determined with enough precision yet). It is worthwhile noting that this method does not depend on the actual measurement of the redshift drift which, as mentioned in the Introduction, is not available yet.

We have shown that the relation given by Eq. (16) can be used to set limits on the parameter space of a given $f(R)$ theory. As examples, we have selected the theory proposed by Hu and Sawicki, and the exponential gravity theory introduced by Linder. For the former, we found a constraint on the possible values of the parameter n that improves previous bound obtained from other cosmological data. In the case of the latter theory, our method ameliorated the previous bound for one of the parameters of the theory (c), and provided the first bounds on the second (r/m^2).

It is worth noting that the approach presented here, which is based on the redshift drift in $f(R)$ theories, can in principle be applied to other observables quantities relevant to cosmology, as well as to other alternative theories of gravity.

To close, we would like to emphasize that the bounds obtained by the method developed here can be analyzed together with those coming from the observations mentioned in the Introduction as well as other means (such as energy conditions [41]), with the aim of deciding whether a given $f(R)$ theory is consistent with the available data.

Acknowledgements

FATP acknowledges support from CONICET and CLAF/ICTP. SEPB would like to acknowledge support from FAPERJ, UERJ, and ICRA Net-Pescara.

-
- [1] M. Li, X.-D. Li, S. Wang, and Y. Wang, *Commun. Theor. Phys.* **56**, 525 (2011), 1103.5870.
 - [2] S. Capozziello and V. Faraoni, *Beyond Einstein Gravity. A survey of Gravitational Theories for Cosmology and Astrophysics* (Springer Science+Business Media, 2011).
 - [3] T. P. Sotiriou and V. Faraoni, *Rev. Mod. Phys.* **82**, 451 (2010), 0805.1726.
 - [4] A. Dolgov and M. Kawasaki, *Phys. Lett. B* **573**, 1 (2003), astro-ph/0307285.
 - [5] S. M. Carroll, M. Hoffman, and M. Trodden, *Phys. Rev. D* **68**, 023509 (2003).
 - [6] L. Amendola, R. Gannouji, D. Polarski, and S. Tsujikawa, *Phys. Rev. D* **75**, 083504 (2007), gr-qc/0612180.
 - [7] A. de Felice and S. Tsujikawa, *Living Reviews in Relativity* **13**, 3 (2010), 1002.4928.
 - [8] S. Nojiri and S. D. Odintsov, *Phys. Rept.* **505**, 59 (2011), 1011.0544.
 - [9] R. Bean, D. Bernat, L. Pogosian, A. Silvestri, and M. Trodden, *Phys. Rev. D* **75**, 064020 (2007).
 - [10] A. A. Starobinsky, *Soviet Journal. of Experimental. and Theoretical. Phys. Lett.* **86**, 157 (2007), 0706.2041.
 - [11] W. Hu and I. Sawicki, *Phys. Rev. D* **76**, 064004 (2007), 0705.1158.
 - [12] S. M. Carroll, I. Sawicki, A. Silvestri, and M. Trodden, *New J. Phys.* **8**, 323 (2006), astro-ph/0607458.
 - [13] A.-C. Davis, E. A. Lim, J. Sakstein, and D. Shaw, *Phys. Rev. D* **85**, 123006 (2012), 1102.5278.
 - [14] P. Brax, S. Clesse, and A.-C. Davis, *JCAP* **1301**, 003 (2013), 1207.1273.
 - [15] J. Lee, G.-B. Zhao, B. Li, and K. Koyama, *Astrophys. J.* **763**, 28 (2013), 1204.6608.
 - [16] V. Cardone, S. Camera, R. Mainini, A. Romano, A. Diaferio, et al. (2012), 1204.3148.
 - [17] A. Sandage, *Astrophys. J.* **136**, 319 (1962).
 - [18] G. C. McVittie, *Astrophys. J.* **136**, 334 (1962).
 - [19] A. Loeb, *Astrophys. Journal Lett.* **499**, L111 (1998), arXiv:astro-ph/9802122.
 - [20] C. Quercellini, L. Amendola, A. Balbi, P. Cabella, and M. Quartin, *Phys. Rept.* **521**, 95 (2012), 1011.2646.
 - [21] C. Quercellini, M. Quartin, and L. Amendola, *Phys. Rev. Lett.* **102**, 151302 (2009).
 - [22] J.-P. Uzan, C. Clarkson, and G. F. Ellis, *Phys. Rev. Lett.* **100**, 191303 (2008), 0801.0068.
 - [23] M. Quartin and L. Amendola, *Phys. Rev. D* **81**, 043522 (2010), 0909.4954.
 - [24] J. Plebanski and A. Krasinski, *An Introduction to General Relativity and Cosmology* (2006).
 - [25] C.-M. Yoo, T. Kai, and K.-I. Nakao, *Phys. Rev. D* **83**, 043527 (2011), 1010.0091.
 - [26] P. E. Vielzeuf and C. J. A. P. Martins, *Phys. Rev. D* **85**, 087301 (2012), 1202.4364.
 - [27] M. Martinelli, S. Pandolfi, C. J. A. P. Martins, and P. E. Vielzeuf, *Phys. Rev. D* **86**, 123001 (2012), 1210.7166.
 - [28] B. Moraes and D. Polarski, *Phys. Rev. D* **84**, 104003 (2011), 1110.2525.
 - [29] S. Weinberg, *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity* (1972).
 - [30] S. Capozziello, V. Cardone, and V. Salzano, *Phys. Rev. D* **78**, 063504 (2008), 0802.1583.
 - [31] S. Capozziello, R. Lazkoz, and V. Salzano, *Phys. Rev. D* **84**, 124061 (2011), 1104.3096.
 - [32] A. Aviles, A. Bravetti, S. Capozziello, and O. Luongo, *ArXiv e-prints* (2012), 1210.5149.
 - [33] A. Shafieloo, A. G. Kim, and E. V. Linder, *Phys. Rev. D* **85**, 123530 (2012), 1204.2272.
 - [34] E. V. Linder, *Phys. Rev. D* **80**, 123528 (2009), 0905.2962.
 - [35] M. Visser, *Gen. Rel. & Grav.* **37**, 1541 (2005).
 - [36] R. Kerner, *Gen. Rel. & Grav.* **14**, 453 (1982).
 - [37] E. Komatsu, K. M. Smith, J. Dunkley, C. L. Bennett, B. Gold, G. Hinshaw, N. Jarosik, D. Larson, M. R.olta, L. Page, et al., *Astrophys. Journal Supp. Series* **192**, 18 (2011), 1001.4538.
 - [38] A. G. Riess, L. Macri, S. Casertano, M. Sosey, H. Lampeitl, H. C. Ferguson, A. V. Filippenko, S. W. Jha, W. Li, R. Chornock, et al., *Astrophys. J.* **699**, 539 (2009), 0905.0695.
 - [39] M. Martinelli, A. Melchiorri, O. Mena, V. Salvatelli, and Z. Gironés, *Phys. Rev. D* **85**, 024006 (2012), 1109.4736.
 - [40] L. Yang, C.-C. Lee, L.-W. Luo, and C.-Q. Geng, *Phys. Rev. D* **82**, 103515 (2010), 1010.2058.
 - [41] S. E. Perez Bergliaffa, *Phys.Lett.* **B642**, 311 (2006), gr-qc/0608072.